

# UP Board Class 12 Mathematics 2026 (Set 324 DD) Question Paper with Solutions



## General Instructions

- (i) This booklet contains 34 questions, each provided with a complete, step-by-step solution.
- (ii) It comprises 5 single-correct multiple-choice questions.
- (iii) Attempt each question on your own before reviewing the given solution.

1. If the function  $f : N \rightarrow N$  is defined as  $f(x) = x^2$ , then  $f$  is:

- (A) One-one and onto
- (B) One-one but not onto
- (C) Neither one-one nor onto
- (D) Many-one and onto

**Correct Answer:** (B) One-one but not onto

### Solution:

#### Step 1: Check one-one:

For  $x_1, x_2 \in N$ , if  $f(x_1) = f(x_2)$  then  $x_1^2 = x_2^2$ . Since  $x_1, x_2$  are natural numbers (positive), this forces  $x_1 = x_2$ . So  $f$  is one-one.

#### Step 2: Check onto:

For  $f$  to be onto, every  $y \in N$  must have a pre-image  $x \in N$  with  $x^2 = y$ , i.e.  $x = \sqrt{y}$ . Take  $y = 2$ :  $\sqrt{2}$  is not a natural number, so 2 has no pre-image in  $N$ .

#### Step 3: Why the other options are wrong:

Option A and D claim  $f$  is onto, which fails as shown. Option C claims  $f$  is not one-one, which is false since we proved injectivity above.

**Final Answer:**

$f$  is one-one but not onto.

One-one but not onto

**Quick Tip:** Squaring is injective on naturals but its range misses non-perfect-squares like 2.

2. The relation  $R = \{(a, b) : a \leq b^2\}$  is defined in the set of real numbers. Then  $R$  is:

- (A) Reflexive and symmetric but not transitive
- (B) Reflexive and transitive but not symmetric
- (C) Symmetric and transitive but not reflexive
- (D) Not reflexive, not symmetric and not transitive

**Correct Answer:** (D) Not reflexive, not symmetric and not transitive

**Solution:**

**Step 1: Check reflexivity:**

Reflexive needs  $a \leq a^2$  for every real  $a$ , i.e.  $a^2 - a \geq 0$ , i.e.  $a(a - 1) \geq 0$ , true only for  $a \leq 0$  or  $a \geq 1$ . Take  $a = 0.5$ :  $0.5 \leq 0.25$  is false. So  $R$  is not reflexive.

**Step 2: Check symmetry:**

Take  $a = 0$ ,  $b = 10$ :  $0 \leq 10^2 = 100$  is true, so  $(0, 10) \in R$ . But is  $(10, 0)$

$\in R$ ? That needs  $10 \leq 0^2 = 0$ , false. So  $R$  is not symmetric.

**Step 3: Check transitivity:**

Take  $a = 10$ ,  $b = 4$ ,  $c = 2$ .  $a \leq b^2$ :  $10 \leq 16$  true.  $b \leq c^2$ :  $4 \leq 4$  true. But  $a \leq c^2$ :  $10 \leq 4$  is false. So  $R$  is not transitive.

**Final Answer:**

$R$  fails reflexivity, symmetry and transitivity all three.

Not reflexive, not symmetric, not transitive

**Quick Tip:** Test  $a=0.5$  for reflexivity,  $(0,10)$  vs  $(10,0)$  for symmetry, and  $(10,4,2)$  for transitivity — all fail.

3. If vectors  $5\hat{i} - \lambda\hat{j} + 2\hat{k}$  and  $2\hat{i} + 3\hat{j} + 4\hat{k}$  are perpendicular to each other, then the value of  $\lambda$  is:

- (A) 3
- (B) 4
- (C) 6
- (D) 0

**Correct Answer:** (C) 6

**Solution:**

**Step 1: Use the perpendicularity condition:**

Two vectors are perpendicular exactly when their dot product is 0.

**Step 2: Compute the dot product:**

$$(5)(2) + (-\lambda)(3) + (2)(4) = 10 - 3\lambda + 8 = 18 - 3\lambda$$

**Step 3: Solve for  $\lambda$ :**

$$18 - 3\lambda = 0 \implies \lambda = 6$$

**Final Answer:**

$$\boxed{\lambda = 6}$$

**Quick Tip:** Set the dot product of the two vectors to zero and solve for  $\lambda$ .



4. If  $x - y = \pi$ , then the value of  $\frac{dy}{dx}$  is:

- (A) 0
- (B)  $\pi$
- (C) 1
- (D)  $x$

**Correct Answer:** (C) 1

**Solution:**

**Step 1: Differentiate both sides w.r.t.  $x$ :**

$x - y = \pi$  is an equation between  $x$  and  $y$  with  $\pi$  a constant.

Differentiating:  $1 - \frac{dy}{dx} = 0$ .

**Step 2:** Solve for  $dy/dx$ :

$$\frac{dy}{dx} = 1$$

**Final Answer:**

$$\boxed{\frac{dy}{dx} = 1}$$

**Quick Tip:** Differentiate  $x - y = \text{constant}$  directly; the derivative of a constant is 0.



**5.** Direction cosines of the  $x$ -axis are:

- (A) (1, 0, 0)
- (B) (0, 1, 0)
- (C) (0, 0, 1)
- (D) (0, 0, 0)

**Correct Answer:** (A) (1, 0, 0)

**Solution:**

**Step 1:** Recall the definition:

Direction cosines are the cosines of the angles the axis/line makes with the  $x$ ,  $y$  and  $z$  axes respectively.

**Step 2:** Apply it to the  $x$ -axis itself:

The  $x$ -axis makes angle  $0^\circ$  with itself, and  $90^\circ$  with both the  $y$ -axis and the  $z$ -axis. So its direction cosines are  $(\cos 0^\circ, \cos 90^\circ, \cos 90^\circ)$   
 $= (1, 0, 0)$ .

**Final Answer:**

$$(1, 0, 0)$$

**Quick Tip:** The  $x$ -axis makes 0 degrees with itself and 90 degrees with the  $y$ - and  $z$ -axes.

6. Find the principal value of  $\sin^{-1}\left(\frac{1}{\sqrt{2}}\right)$ .

**Correct Answer:** —

**Solution:**

**Step 1: Set up the equation:**

Let  $\theta = \sin^{-1}\left(\frac{1}{\sqrt{2}}\right)$ , so  $\sin \theta = \frac{1}{\sqrt{2}}$ , with  $\theta \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$  (the principal range).

**Step 2: Identify the angle:**

$\sin\left(\frac{\pi}{4}\right) = \frac{1}{\sqrt{2}}$ , and  $\frac{\pi}{4}$  lies in the principal range.

**Final Answer:**

$$\theta = \frac{\pi}{4}$$

**Quick Tip:** Recall  $\sin(\pi/4) = 1/\sqrt{2}$ , and  $\pi/4$  lies in the principal range of  $\sin^{-1}$ .

7. Find the value of  $\int_2^3 \frac{1}{x} dx$ .

**Correct Answer:** —

**Solution:**

**Step 1: Find the antiderivative:**

$$\int \frac{1}{x} dx = \ln|x| + C.$$

**Step 2: Apply the limits:**

$$\int_2^3 \frac{1}{x} dx = \ln 3 - \ln 2 = \ln \frac{3}{2}$$

**Final Answer:**

$$\boxed{\ln \frac{3}{2}}$$

**Quick Tip:** The antiderivative of  $1/x$  is  $\ln|x|$ ; evaluate at the limits and subtract.

8. Find the angle between vectors  $\hat{i} - 2\hat{j} + 3\hat{k}$  and  $3\hat{i} - 2\hat{j} + \hat{k}$ .

**Correct Answer:** —

### Solution:

**Step 1:** Compute the dot product:

$$(1)(3) + (-2)(-2) + (3)(1) = 3 + 4 + 3 = 10$$

**Step 2:** Compute the magnitudes:

$$|\vec{a}| = \sqrt{1^2 + (-2)^2 + 3^2} = \sqrt{14}, \quad |\vec{b}| = \sqrt{3^2 + (-2)^2 + 1^2} = \sqrt{14}$$

**Step 3:** Use  $\cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}||\vec{b}|}$ :

$$\cos \theta = \frac{10}{\sqrt{14} \cdot \sqrt{14}} = \frac{10}{14} = \frac{5}{7}$$

**Final Answer:**

$$\theta = \cos^{-1}\left(\frac{5}{7}\right)$$

**Quick Tip:** Use  $\cos(\theta) = (a \cdot b) / (|a||b|)$  with the dot product and magnitudes of the two vectors.

9. Direction ratios of a line are 2, -1 and -2. Find the direction cosines of the line.

**Correct Answer:** —

**Solution:**

**Step 1:** Find the magnitude of the direction ratio vector:

$$\sqrt{2^2 + (-1)^2 + (-2)^2} = \sqrt{4 + 1 + 4} = \sqrt{9} = 3$$

**Step 2:** Divide each ratio by the magnitude:

$$\left(\frac{2}{3}, -\frac{1}{3}, -\frac{2}{3}\right)$$

**Final Answer:**

$$\boxed{\left(\frac{2}{3}, -\frac{1}{3}, -\frac{2}{3}\right)}$$

**Quick Tip:** Divide each direction ratio by the square root of the sum of their squares.

10. If  $P(A) = \frac{7}{13}$ ,  $P(B) = \frac{9}{13}$  and  $P(A \cap B) = \frac{4}{13}$ , then find  $P\left(\frac{A}{B}\right)$ .

**Correct Answer:** —

**Solution:**

**Step 1:** Recall the conditional probability formula:

$$P\left(\frac{A}{B}\right) = \frac{P(A \cap B)}{P(B)}$$

**Step 2:** Substitute the given values:

$$P\left(\frac{A}{B}\right) = \frac{4/13}{9/13} = \frac{4}{9}$$

**Final Answer:**

$$\boxed{\frac{4}{9}}$$

**Quick Tip:** Use  $P(A|B) = P(A \text{ intersection } B) / P(B)$ ; the 13-denominators cancel.

**11.** If functions  $f : R \rightarrow R$  and  $g : R \rightarrow R$  are defined respectively as  $f(x) = \cos x$  and  $g(x) = 3x^2$ , then find  $gof$  and  $fog$ .

**Correct Answer:** —

**Solution:**

**Step 1:** Compute  $gof(x) = g(f(x))$ :

$$f(x) = \cos x, \text{ so } g(f(x)) = g(\cos x) = 3(\cos x)^2 = 3 \cos^2 x.$$

**Step 2:** Compute  $fog(x) = f(g(x))$ :

$$g(x) = 3x^2, \text{ so } f(g(x)) = f(3x^2) = \cos(3x^2).$$

**Final Answer:**

$$\boxed{gof(x) = 3 \cos^2 x, \quad fog(x) = \cos(3x^2)}$$

**Quick Tip:** Substitute  $f$  into  $g$  for  $gof$ , and  $g$  into  $f$  for  $fog$ , in that order.

12. Find the value of the determinant  $\begin{vmatrix} x & a & x+a \\ y & b & y+b \\ z & c & z+c \end{vmatrix}$ .

**Correct Answer:** —

**Solution:**

**Step 1:** Compare the third column to the first two:

In every row, the third entry is the sum of the first two:  $x + a$ ,  $y + b$ ,  $z + c$ . So column 3 = column 1 + column 2.

**Step 2:** Apply the column operation  $C_3 \rightarrow C_3 - C_1 - C_2$ :

This is a legal determinant operation (adding a multiple of other columns doesn't change the value) and turns column 3 entirely to zeros.

**Step 3:** A determinant with a zero column is zero:

Expanding along the (now all-zero) third column gives 0.

**Final Answer:**

$$\boxed{0}$$

**Quick Tip:** Column 3 is exactly column 1 + column 2 in every row, so the columns are linearly dependent.

13. If  $f(x) = \begin{cases} x + 2, & x \neq 0 \\ 1, & x = 0 \end{cases}$ , then prove that the function is not continuous at  $x = 0$ .

**Correct Answer:** —

**Solution:**

**Step 1: Find the limit as  $x \rightarrow 0$ :**

For  $x \neq 0$ ,  $f(x) = x + 2$ , so  $\lim_{x \rightarrow 0} f(x) = 0 + 2 = 2$ .

**Step 2: Find the function's actual value at  $x = 0$ :**

By definition,  $f(0) = 1$ .

**Step 3: Compare:**

Continuity at  $x = 0$  requires  $\lim_{x \rightarrow 0} f(x) = f(0)$ . Here  $2 \neq 1$ .

**Final Answer:**

Since the limit and the function value disagree,  $f$  is not continuous at  $x = 0$ .

$$\boxed{\lim_{x \rightarrow 0} f(x) = 2 \neq f(0) = 1}$$

**Quick Tip:** Compute the limit of  $x+2$  as  $x$  approaches 0 and compare it to the defined value  $f(0)=1$ .

14. Find the vector and Cartesian equations of a line which passes through the point  $(1, 2, 3)$  and is parallel to the vector  $2\hat{i} + 3\hat{j} + 2\hat{k}$ .

**Correct Answer:** —

**Solution:**

**Step 1: Write the vector equation:**

A line through point  $\vec{a} = \hat{i} + 2\hat{j} + 3\hat{k}$  parallel to  $\vec{b} = 2\hat{i} + 3\hat{j} + 2\hat{k}$  has vector equation  $\vec{r} = \vec{a} + \lambda\vec{b}$ .

$$\vec{r} = (\hat{i} + 2\hat{j} + 3\hat{k}) + \lambda(2\hat{i} + 3\hat{j} + 2\hat{k})$$

**Step 2: Convert to Cartesian form:**

With  $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$ , matching components gives  $x = 1 + 2\lambda$ ,  $y = 2 + 3\lambda$ ,  $z = 3 + 2\lambda$ , so

$$\frac{x-1}{2} = \frac{y-2}{3} = \frac{z-3}{2}$$

**Final Answer:**

$$\vec{r} = (\hat{i} + 2\hat{j} + 3\hat{k}) + \lambda(2\hat{i} + 3\hat{j} + 2\hat{k}), \quad \frac{x-1}{2} = \frac{y-2}{3} = \frac{z-3}{2}$$

**Quick Tip:** Use  $r = a + \lambda b$  for the vector form; equate components for the Cartesian symmetric form.

15. Find the general solution of the differential equation  $\frac{dy}{dx} = \frac{1+y^2}{1+x^2}$ .

**Correct Answer:** —

**Solution:**

**Step 1: Separate the variables:**

$$\frac{dy}{1+y^2} = \frac{dx}{1+x^2}$$

**Step 2: Integrate both sides:**

$$\int \frac{dy}{1+y^2} = \int \frac{dx}{1+x^2}$$

$$\tan^{-1} y = \tan^{-1} x + C$$

**Final Answer:**

$$\boxed{\tan^{-1} y - \tan^{-1} x = C}$$

**Quick Tip:** This is variable-separable; both sides integrate to an arctangent.



16. If  $y = 5 \cos x - 3 \sin x$ , then prove that  $\frac{d^2y}{dx^2} + y = 0$ .

**Correct Answer:** —

**Solution:**

**Step 1: Differentiate once:**

$$\frac{dy}{dx} = -5 \sin x - 3 \cos x$$

**Step 2: Differentiate again:**

$$\frac{d^2y}{dx^2} = -5 \cos x + 3 \sin x = -(5 \cos x - 3 \sin x) = -y$$

**Step 3: Rearrange:**

$$\frac{d^2y}{dx^2} + y = -y + y = 0$$

**Final Answer:**

$$\boxed{\frac{d^2y}{dx^2} + y = 0}$$

**Quick Tip:** Differentiate twice and notice the second derivative is exactly -y.

17. Find the projection of vector  $\hat{i} + 3\hat{j} + 7\hat{k}$  on vector  $7\hat{i} - \hat{j} + 8\hat{k}$ .

**Correct Answer:** —

**Solution:**

**Step 1:** Recall the projection formula:

Projection of  $\vec{a}$  on  $\vec{b}$  is  $\frac{\vec{a} \cdot \vec{b}}{|\vec{b}|}$ .

**Step 2:** Compute the dot product:

$$(1)(7) + (3)(-1) + (7)(8) = 7 - 3 + 56 = 60$$

**Step 3:** Compute  $|\vec{b}|$ :

$$|\vec{b}| = \sqrt{7^2 + (-1)^2 + 8^2} = \sqrt{49 + 1 + 64} = \sqrt{114}$$

**Final Answer:**

$$\frac{60}{\sqrt{114}}$$

**Quick Tip:** Projection of  $a$  on  $b$  equals  $(a \cdot b)/|b|$ ; compute the dot product and the magnitude of  $b$ .

18. If  $P(A) = \frac{1}{3}$ ,  $P(B) = \frac{1}{2}$  and  $P(A \cup B) = \frac{2}{3}$ , then show that  $A$  and  $B$  are independent events.

**Correct Answer:** —

**Solution:**

**Step 1:** Find  $P(A \cap B)$  using the addition rule:

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$\frac{2}{3} = \frac{1}{3} + \frac{1}{2} - P(A \cap B) \implies P(A \cap B) = \frac{1}{3} + \frac{1}{2} - \frac{2}{3} = \frac{2+3-4}{6} = \frac{1}{6}$$

**Step 2:** Compute  $P(A) \cdot P(B)$ :

$$P(A) \cdot P(B) = \frac{1}{3} \times \frac{1}{2} = \frac{1}{6}$$

**Step 3:** Compare:

$P(A \cap B) = \frac{1}{6} = P(A) \cdot P(B)$ , which is exactly the independence condition.

**Final Answer:**

Since  $P(A \cap B) = P(A)P(B)$ ,  $A$  and  $B$  are independent.

$$P(A \cap B) = P(A)P(B) = \frac{1}{6}$$

**Quick Tip:** Find  $P(A \text{ intersect } B)$  from the addition rule, then check if it equals  $P(A)$  times  $P(B)$ .

19. Prove that  $\tan^{-1} \frac{1}{5} + \tan^{-1} \frac{1}{7} + \tan^{-1} \frac{1}{3} + \tan^{-1} \frac{1}{8} = \frac{\pi}{4}$ .

**Correct Answer:** —

### Solution:

**Step 1:** Combine the first pair using  $\tan^{-1} x + \tan^{-1} y = \tan^{-1} \frac{x+y}{1-xy}$ :

$$\tan^{-1} \frac{1}{5} + \tan^{-1} \frac{1}{7} = \tan^{-1} \frac{\frac{1}{5} + \frac{1}{7}}{1 - \frac{1}{5} \cdot \frac{1}{7}} = \tan^{-1} \frac{12/35}{34/35} = \tan^{-1} \frac{12}{34} = \tan^{-1} \frac{6}{17}$$

**Step 2:** Combine the second pair the same way:

$$\tan^{-1} \frac{1}{3} + \tan^{-1} \frac{1}{8} = \tan^{-1} \frac{\frac{1}{3} + \frac{1}{8}}{1 - \frac{1}{3} \cdot \frac{1}{8}} = \tan^{-1} \frac{11/24}{23/24} = \tan^{-1} \frac{11}{23}$$

**Step 3:** Combine the two results:

$$\tan^{-1} \frac{6}{17} + \tan^{-1} \frac{11}{23} = \tan^{-1} \frac{\frac{6}{17} + \frac{11}{23}}{1 - \frac{6}{17} \cdot \frac{11}{23}}$$

$$\text{Numerator: } \frac{6 \times 23 + 11 \times 17}{17 \times 23} = \frac{138 + 187}{391} = \frac{325}{391}.$$

$$\text{Denominator: } 1 - \frac{66}{391} = \frac{325}{391}.$$

$$\text{So the fraction is } \frac{325/391}{325/391} = 1.$$

**Step 4:** Conclude:

$$\tan^{-1}(1) = \frac{\pi}{4}$$

**Final Answer:**

$$\tan^{-1} \frac{1}{5} + \tan^{-1} \frac{1}{7} + \tan^{-1} \frac{1}{3} + \tan^{-1} \frac{1}{8} = \frac{\pi}{4}$$

**Quick Tip:** Pair the terms and repeatedly apply the tan inverse addition formula; the final ratio collapses to 1.

**20.** If  $\cos y = x \cos(a + y)$  and  $\cos a \neq \pm 1$ , then prove that  $\frac{dy}{dx} = \frac{\cos^2(a + y)}{\sin a}$ .

**Correct Answer:** —

**Solution:**

**Step 1:** Differentiate both sides implicitly w.r.t.  $x$ :

$$-\sin y \cdot \frac{dy}{dx} = \cos(a + y) + x \cdot (-\sin(a + y)) \frac{dy}{dx}$$

**Step 2:** Collect the  $dy/dx$  terms:

$$-\sin y \cdot y' + x \sin(a + y) \cdot y' = \cos(a + y)$$

$$y' [x \sin(a + y) - \sin y] = \cos(a + y)$$

**Step 3:** Substitute  $x = \frac{\cos y}{\cos(a + y)}$  from the original equation:

$$x \sin(a + y) - \sin y = \frac{\cos y \sin(a + y)}{\cos(a + y)} - \sin y = \frac{\cos y \sin(a + y) - \sin y \cos(a + y)}{\cos(a + y)}$$

The numerator is  $\sin[(a + y) - y] = \sin a$  (sine subtraction formula), so this simplifies to  $\frac{\sin a}{\cos(a + y)}$ .

**Step 4: Solve for  $y'$ :**

$$y' \cdot \frac{\sin a}{\cos(a + y)} = \cos(a + y) \implies y' = \frac{\cos^2(a + y)}{\sin a}$$

**Final Answer:**

$$\boxed{\frac{dy}{dx} = \frac{\cos^2(a + y)}{\sin a}}$$

**Quick Tip:** Differentiate implicitly, then substitute  $x$  back from the original relation and use  $\sin(A-B)$ .

**21.** Solve the differential equation:  $(1 + x^2) dy + 2xy dx = \cot x dx$ .

**Correct Answer:** —

**Solution:**

**Step 1: Write in standard linear form  $\frac{dy}{dx} + Py = Q$ :**

Divide throughout by  $dx$  and by  $(1 + x^2)$ :

$$\frac{dy}{dx} + \frac{2x}{1 + x^2} y = \frac{\cot x}{1 + x^2}$$

Here  $P = \frac{2x}{1+x^2}$ ,  $Q = \frac{\cot x}{1+x^2}$ .

**Step 2: Find the integrating factor:**

$$\text{IF} = e^{\int P dx} = e^{\int \frac{2x}{1+x^2} dx} = e^{\ln(1+x^2)} = 1+x^2$$

**Step 3: Multiply through by the IF and integrate:**

$$\frac{d}{dx} [y(1+x^2)] = \cot x$$

$$y(1+x^2) = \int \cot x dx = \ln |\sin x| + C$$

**Final Answer:**

$$y(1+x^2) = \ln |\sin x| + C$$

**Quick Tip:** Convert to linear form  $dy/dx + Py = Q$ , find  $\text{IF} = 1+x^2$ , and integrate  $\cot x$ .

**22.** Maximize  $Z = 10x + 3y$  by the graphical method under the following constraints:  $x \geq 0$ ;  $y \geq 0$ ;  $5x + 3y \leq 15$ ;  $2x + 5y \leq 10$ .

**Correct Answer:** —

## Solution:

### Step 1: Find the feasible-region corner points:

The boundary lines are  $5x + 3y = 15$  and  $2x + 5y = 10$ , together with the axes. Intercepts:  $5x + 3y = 15$  meets axes at  $(3, 0)$  and  $(0, 5)$ ;  $2x + 5y = 10$  meets axes at  $(5, 0)$  and  $(0, 2)$ .

### Step 2: Check which intercepts are feasible (satisfy both constraints):

$(3, 0)$ : check  $2(3) + 5(0) = 6 \leq 10$  ✓ — feasible.

$(0, 2)$ : check  $5(0) + 3(2) = 6 \leq 15$  ✓ — feasible.

$(0, 5)$ : check  $2(0) + 5(5) = 25 \leq 10$ ? No — infeasible.

$(5, 0)$ : check  $5(5) + 3(0) = 25 \leq 15$ ? No — infeasible.

### Step 3: Find the intersection of the two lines:

From  $5x + 3y = 15$ :  $x = 3 - 0.6y$ . Substitute into  $2x + 5y = 10$ :  $2(3 - 0.6y) + 5y = 10 \Rightarrow 6 - 1.2y + 5y = 10 \Rightarrow 3.8y = 4 \Rightarrow y = \frac{20}{19}$ ,  $x = \frac{45}{19}$ .

### Step 4: Evaluate $Z$ at every feasible corner: $(0, 0)$ , $(3, 0)$ , $(0, 2)$ ,

$(\frac{45}{19}, \frac{20}{19})$ :

$Z(0, 0) = 0$ .  $Z(3, 0) = 30$ .  $Z(0, 2) = 6$ .  $Z(\frac{45}{19}, \frac{20}{19}) = \frac{450 + 60}{19} = \frac{510}{19} \approx 26.8$ .

### Final Answer:

The maximum occurs at  $(3, 0)$ .

$$Z_{\max} = 30 \text{ at } (x, y) = (3, 0)$$

**Quick Tip:** Plot the two constraint lines, find all feasible corner points, and evaluate  $Z$  at each.

**23.** Find the shortest distance between the lines  $\vec{r} = (\hat{i} + 2\hat{j} + \hat{k}) + \lambda(\hat{i} - \hat{j} + \hat{k})$  and  $\vec{r} = (2\hat{i} - \hat{j} - \hat{k}) + \mu(2\hat{i} + \hat{j} + 2\hat{k})$ .

**Correct Answer:** —

**Solution:**

**Step 1:** Identify  $\vec{a}_1, \vec{b}_1, \vec{a}_2, \vec{b}_2$ :

$$\vec{a}_1 = (1, 2, 1), \vec{b}_1 = (1, -1, 1), \vec{a}_2 = (2, -1, -1), \vec{b}_2 = (2, 1, 2).$$

**Step 2:** Compute  $\vec{a}_2 - \vec{a}_1$ :

$$\vec{a}_2 - \vec{a}_1 = (1, -3, -2)$$

**Step 3:** Compute  $\vec{b}_1 \times \vec{b}_2$ :

$$\vec{b}_1 \times \vec{b}_2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -1 & 1 \\ 2 & 1 & 2 \end{vmatrix} = \hat{i}(-2 - 1) - \hat{j}(2 - 2) + \hat{k}(1 + 2) = (-3, 0, 3)$$

**Step 4:** Compute the shortest distance formula:

$$d = \frac{|(\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2)|}{|\vec{b}_1 \times \vec{b}_2|}$$

Numerator:  $(1)(-3) + (-3)(0) + (-2)(3) = -3 + 0 - 6 = -9$ , so  $|-9| = 9$ .

Denominator:  $|(-3, 0, 3)| = \sqrt{9 + 0 + 9} = \sqrt{18} = 3\sqrt{2}$ .

**Final Answer:**

$$d = \frac{9}{3\sqrt{2}} = \frac{3}{\sqrt{2}} = \frac{3\sqrt{2}}{2}$$

$$\boxed{d = \frac{3\sqrt{2}}{2}}$$

**Quick Tip:** Use  $d = |(a_2 - a_1) \cdot (b_1 \times b_2)| / |b_1 \times b_2|$  for two skew lines in vector form.

24. Find the value of the determinant  $\begin{vmatrix} a & b & c \\ a^2 & b^2 & c^2 \\ a^3 & b^3 & c^3 \end{vmatrix}$ .

**Correct Answer:** —

**Solution:**

**Step 1: Factor each column:**

Column 1 is  $a, a^2, a^3 = a(1, a, a^2)$ ; similarly column 2 factors out  $b$ , column 3 factors out  $c$ . Pulling these common factors out of each column:

$$\begin{vmatrix} a & b & c \\ a^2 & b^2 & c^2 \\ a^3 & b^3 & c^3 \end{vmatrix} = abc \begin{vmatrix} 1 & 1 & 1 \\ a & b & c \\ a^2 & b^2 & c^2 \end{vmatrix}$$

**Step 2: Recognize the remaining determinant as a Vandermonde determinant:**

$$\begin{vmatrix} 1 & 1 & 1 \\ a & b & c \\ a^2 & b^2 & c^2 \end{vmatrix} = (b-a)(c-a)(c-b)$$

**Step 3:** Rewrite the sign in the conventional  $(a-b)(b-c)(c-a)$  form:

$(b-a)(c-a)(c-b) = (a-b)(b-c)(c-a)$  (each pair of sign flips cancels — two flips leave the product unchanged).

**Final Answer:**

$$abc(a-b)(b-c)(c-a)$$

**Quick Tip:** Factor  $a, b, c$  out of the columns to expose a Vandermonde determinant, then apply its known factorization.

25. Evaluate  $\int \frac{x^2 + 1}{x^2 - 5x + 6} dx$ .

**Correct Answer:** —

**Solution:**

**Step 1:** Since numerator and denominator have equal degree, do polynomial division:

$$\frac{x^2 + 1}{x^2 - 5x + 6} = 1 + \frac{5x - 5}{x^2 - 5x + 6}$$

**Step 2: Factor the denominator and set up partial fractions:**

$$x^2 - 5x + 6 = (x - 2)(x - 3). \text{ Write } \frac{5x - 5}{(x - 2)(x - 3)} = \frac{A}{x - 2} + \frac{B}{x - 3}.$$

**Step 3: Solve for A and B:**

$$5x - 5 = A(x - 3) + B(x - 2). \text{ At } x = 2: 5 = A(-1) \Rightarrow A = -5. \text{ At } x = 3: 10 = B(1) \Rightarrow B = 10.$$

**Step 4: Integrate term by term:**

$$\int \left[ 1 - \frac{5}{x - 2} + \frac{10}{x - 3} \right] dx = x - 5 \ln |x - 2| + 10 \ln |x - 3| + C$$

**Final Answer:**

$$x - 5 \ln |x - 2| + 10 \ln |x - 3| + C$$

**Quick Tip:** Do polynomial long division first (equal degree top and bottom), then partial fractions on the remainder.

**26.** Prove that vectors  $2\hat{i} - \hat{j} + \hat{k}$ ,  $\hat{i} - 3\hat{j} - 5\hat{k}$  and  $3\hat{i} - 4\hat{j} - 4\hat{k}$  are vertices of a right-angled triangle.

**Correct Answer:** —

**Solution:**

**Step 1: Label the points and find the side vectors:**

$$\text{Let } A = (2, -1, 1), B = (1, -3, -5), C = (3, -4, -4).$$

$$\vec{AB} = B - A = (-1, -2, -6), \vec{BC} = C - B = (2, -1, 1), \vec{CA} = A - C$$

$$= (-1, 3, 5).$$

**Step 2:** Compute the squared lengths of the three sides:

$$|AB|^2 = 1 + 4 + 36 = 41, |BC|^2 = 4 + 1 + 1 = 6, |CA|^2 = 1 + 9 + 25 = 35.$$

**Step 3:** Check the Pythagorean relation:

$|BC|^2 + |CA|^2 = 6 + 35 = 41 = |AB|^2$ . This matches, so the triangle is right-angled with the right angle at the vertex common to sides  $BC$  and  $CA$ , which is  $C$ .

**Final Answer:**

Since  $BC^2 + CA^2 = AB^2$ , the triangle is right-angled at  $C$ .

$$BC^2 + CA^2 = AB^2 = 41$$

**Quick Tip:** Compute the three side-vectors, their squared lengths, and check which pair satisfies the Pythagorean relation.



27. If  $y = Ae^{mx} + Be^{nx}$ , then show that  $\frac{d^2y}{dx^2} - (m+n)\frac{dy}{dx} + mny = 0$ .

**Correct Answer:** —

**Solution:**

**Step 1:** Find the first derivative:

$$y' = Am e^{mx} + Bn e^{nx}$$

**Step 2:** Find the second derivative:

$$y'' = Am^2e^{mx} + Bn^2e^{nx}$$

**Step 3:** Substitute into  $y'' - (m + n)y' + mny$  and group by  $e^{mx}$  and  $e^{nx}$ :

Coefficient of  $Ae^{mx}$ :  $m^2 - (m + n)m + mn = m^2 - m^2 - mn + mn = 0$ .

Coefficient of  $Be^{nx}$ :  $n^2 - (m + n)n + mn = n^2 - mn - n^2 + mn = 0$ .

**Step 4:** Both coefficients vanish, so the whole expression is zero:

$$y'' - (m + n)y' + mny = 0 \cdot Ae^{mx} + 0 \cdot Be^{nx} = 0$$

**Final Answer:**

$$\boxed{\frac{d^2y}{dx^2} - (m + n)\frac{dy}{dx} + mny = 0}$$

**Quick Tip:** Differentiate twice, substitute into the given expression, and show the coefficients of  $e^{mx}$  and  $e^{nx}$  each vanish.

**28.** If a die is thrown three times, then find the probability of getting at least one odd number.

**Correct Answer:** —

**Solution:**

**Step 1:** Find the probability of NOT getting an odd number on a single throw:

A die has 3 odd faces (1,3,5) and 3 even faces (2,4,6), so  $P(\text{even}) = \frac{3}{6} = \frac{1}{2}$ .

**Step 2:** Find the probability of getting NO odd number in all three throws:

The three throws are independent, so  $P(\text{no odd in 3 throws}) = \left(\frac{1}{2}\right)^3 = \frac{1}{8}$ .

**Step 3:** Use the complement rule:

$$P(\text{at least one odd}) = 1 - P(\text{no odd at all}) = 1 - \frac{1}{8} = \frac{7}{8}$$

**Final Answer:**

$$\boxed{\frac{7}{8}}$$

**Quick Tip:** Use the complement:  $1 - P(\text{all three throws are even})$ .

29. If  $A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 4 & 5 \\ 3 & 5 & 6 \end{bmatrix}$ , then find  $A^{-1}$ .

**Correct Answer:** —

**Solution:**

**Step 1:** Find  $|A|$  by expanding along the first row:

$$|A| = 1(4 \cdot 6 - 5 \cdot 5) - 2(2 \cdot 6 - 5 \cdot 3) + 3(2 \cdot 5 - 4 \cdot 3)$$

$$= 1(24 - 25) - 2(12 - 15) + 3(10 - 12) = -1 + 6 - 6 = -1$$

**Step 2: Find all nine cofactors:**

$$C_{11} = 24 - 25 = -1, C_{12} = -(12 - 15) = 3, C_{13} = 10 - 12 = -2$$

$$C_{21} = -(12 - 15) = 3, C_{22} = 6 - 9 = -3, C_{23} = -(5 - 6) = 1$$

$$C_{31} = 10 - 12 = -2, C_{32} = -(5 - 6) = 1, C_{33} = 4 - 4 = 0$$

**Step 3: Build the adjugate (transpose of the cofactor matrix):**

$$\text{adj}(A) = \begin{bmatrix} -1 & 3 & -2 \\ 3 & -3 & 1 \\ -2 & 1 & 0 \end{bmatrix}$$

**Step 4: Divide by  $|A| = -1$ :**

$$A^{-1} = \frac{1}{-1} \begin{bmatrix} -1 & 3 & -2 \\ 3 & -3 & 1 \\ -2 & 1 & 0 \end{bmatrix} = \begin{bmatrix} 1 & -3 & 2 \\ -3 & 3 & -1 \\ 2 & -1 & 0 \end{bmatrix}$$

**Final Answer:**

$$A^{-1} = \begin{bmatrix} 1 & -3 & 2 \\ -3 & 3 & -1 \\ 2 & -1 & 0 \end{bmatrix}$$

**Quick Tip:** Compute  $|A|$  via cofactor expansion, build the adjugate from all nine cofactors, then divide by  $|A|$ .



**30.** Solve the following system of equations by using the Matrix Method:

$$x - y + 2z = 7$$

$$3x + 4y - 5z = -5$$

$$2x - y + 3z = 12$$

**Correct Answer:** —

**Solution:**

**Step 1:** Write as  $AX = B$  and find  $|A|$ :

$$A = \begin{bmatrix} 1 & -1 & 2 \\ 3 & 4 & -5 \\ 2 & -1 & 3 \end{bmatrix}, B = \begin{bmatrix} 7 \\ -5 \\ 12 \end{bmatrix}.$$

$$|A| = 1(12 - 5) - (-1)(9 + 10) + 2(-3 - 8) = 7 + 19 - 22 = 4$$

**Step 2:** Compute all nine cofactors:

$$C_{11} = 12 - 5 = 7, C_{12} = -(9 + 10) = -19, C_{13} = -3 - 8 = -11$$

$$C_{21} = -(-3 + 2) = 1, C_{22} = 3 - 4 = -1, C_{23} = -(-1 + 2) = -1$$

$$C_{31} = 5 - 8 = -3, C_{32} = -(-5 - 6) = 11, C_{33} = 4 + 3 = 7$$

**Step 3:** Build the adjugate and divide by  $|A| = 4$  to get  $A^{-1}$ :

$$\text{adj}(A) = \begin{bmatrix} 7 & 1 & -3 \\ -19 & -1 & 11 \\ -11 & -1 & 7 \end{bmatrix}, \quad A^{-1} = \frac{1}{4} \begin{bmatrix} 7 & 1 & -3 \\ -19 & -1 & 11 \\ -11 & -1 & 7 \end{bmatrix}$$

**Step 4: Compute  $X = A^{-1}B$ :**

$\text{adj}(A) \cdot B$ : row1 =  $7(7) + 1(-5) - 3(12) = 49 - 5 - 36 = 8$ ; row2 =  $-19(7) - 1(-5) + 11(12) = -133 + 5 + 132 = 4$ ; row3 =  $-11(7) - 1(-5) + 7(12) = -77 + 5 + 84 = 12$ .

$$X = \frac{1}{4} \begin{bmatrix} 8 \\ 4 \\ 12 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix}$$

**Final Answer:**

$$x = 2, y = 1, z = 3$$

**Quick Tip:** Find  $A^{-1}$  via adjugate/determinant, then compute  $X = A^{-1}B$ .

**31.** Prove that the height of the cylinder of maximum volume inscribed in a sphere of radius  $R$  is  $\frac{2R}{\sqrt{3}}$ . Also find the maximum volume of the cylinder.

**Correct Answer:** —

**Solution:**

**Step 1: Set up the constraint between cylinder half-height and radius:**

Let the cylinder have height  $H = 2h$  and base radius  $r$ . Since the cylinder is inscribed in a sphere of radius  $R$ , the center of the sphere, the center of the cylinder's axis and a top-rim point form a right triangle:  $r^2 + h^2 = R^2$ .

**Step 2: Write the volume as a function of  $h$  alone:**

$$V = \pi r^2 H = \pi(R^2 - h^2)(2h) = 2\pi(R^2 h - h^3)$$

**Step 3: Differentiate and set to zero:**

$$\frac{dV}{dh} = 2\pi(R^2 - 3h^2) = 0 \implies h^2 = \frac{R^2}{3} \implies h = \frac{R}{\sqrt{3}}$$

Check it's a maximum:  $\frac{d^2V}{dh^2} = 2\pi(-6h) < 0$  for  $h > 0$ , confirming a maximum.

**Step 4: Find the height and the maximum volume:**

$$H = 2h = \frac{2R}{\sqrt{3}} \quad (\text{as required})$$

$$r^2 = R^2 - h^2 = R^2 - \frac{R^2}{3} = \frac{2R^2}{3}$$

$$V_{\max} = \pi r^2 H = \pi \cdot \frac{2R^2}{3} \cdot \frac{2R}{\sqrt{3}} = \frac{4\pi R^3}{3\sqrt{3}} = \frac{4\sqrt{3}\pi R^3}{9}$$

**Final Answer:**

$$H = \frac{2R}{\sqrt{3}}, \quad V_{max} = \frac{4\pi R^3}{3\sqrt{3}}$$

**Quick Tip:** Express  $V$  as a function of the half-height  $h$  using  $r^2 + h^2 = R^2$ , then maximize via  $dV/dh = 0$ .

**32.** Solve the differential equation  $(x - y)\frac{dy}{dx} = x + 2y$ .

**Correct Answer:** —

**Solution:**

**Step 1:** Recognize this as a homogeneous equation and substitute  $y = vx$ :

$\frac{dy}{dx} = v + x\frac{dv}{dx}$ . The equation  $(x - y)\frac{dy}{dx} = x + 2y$  becomes  $x(1 - v)\left(v + x\frac{dv}{dx}\right) = x(1 + 2v)$ , i.e.  $(1 - v)\left(v + x\frac{dv}{dx}\right) = 1 + 2v$ .

**Step 2:** Expand and isolate the  $x\frac{dv}{dx}$  term:

$$v - v^2 + x(1 - v)\frac{dv}{dx} = 1 + 2v$$

$$x(1 - v)\frac{dv}{dx} = 1 + 2v - v + v^2 = v^2 + v + 1$$

**Step 3: Separate variables:**

$$\frac{1-v}{v^2+v+1} dv = \frac{dx}{x}$$

**Step 4: Integrate the left side by splitting  $1-v = -\frac{1}{2}(2v+1) + \frac{3}{2}$ :**

$$\int \frac{1-v}{v^2+v+1} dv = -\frac{1}{2} \ln(v^2+v+1) + \sqrt{3} \tan^{-1}\left(\frac{2v+1}{\sqrt{3}}\right)$$

(using  $\int \frac{2v+1}{v^2+v+1} dv = \ln(v^2+v+1)$  and completing the square  $v^2+v+1 = (v+\frac{1}{2})^2 + \frac{3}{4}$ ).

**Step 5: Equate to  $\ln|x| + C$  and substitute back  $v = y/x$ :**

$$-\frac{1}{2} \ln\left(\frac{x^2+xy+y^2}{x^2}\right) + \sqrt{3} \tan^{-1}\left(\frac{x+2y}{x\sqrt{3}}\right) = \ln|x| + C$$

The  $\ln x^2$  term combines with  $\ln|x|$  on the right and simplifies to:

**Final Answer:**

$$\sqrt{3} \tan^{-1}\left(\frac{x+2y}{\sqrt{3}x}\right) - \frac{1}{2} \ln(x^2+xy+y^2) = C$$

**Quick Tip:** Substitute  $y=vx$  (homogeneous equation), separate variables, and complete the square in the denominator.

33. Find the value of:  $\int_0^{\pi/2} \log \sin x \, dx$ .

**Correct Answer:** —

**Solution:**

**Step 1:** Let  $I = \int_0^{\pi/2} \log \sin x \, dx$  and use the property  $\int_0^a f(x) dx = \int_0^a f(a-x) dx$ :

$$I = \int_0^{\pi/2} \log \sin \left( \frac{\pi}{2} - x \right) dx = \int_0^{\pi/2} \log \cos x \, dx$$

**Step 2:** Add the two expressions for  $I$ :

$$2I = \int_0^{\pi/2} (\log \sin x + \log \cos x) dx = \int_0^{\pi/2} \log(\sin x \cos x) dx = \int_0^{\pi/2} \log \left( \frac{\sin 2x}{2} \right) dx$$

$$2I = \int_0^{\pi/2} \log \sin 2x \, dx - \int_0^{\pi/2} \log 2 \, dx$$

**Step 3:** Evaluate  $\int_0^{\pi/2} \log \sin 2x \, dx$  via the substitution  $t = 2x$ :

$$\int_0^{\pi/2} \log \sin 2x \, dx = \frac{1}{2} \int_0^{\pi} \log \sin t \, dt = \frac{1}{2} \cdot 2 \int_0^{\pi/2} \log \sin t \, dt = I$$

(using symmetry of  $\sin t$  about  $t = \pi/2$  on  $[0, \pi]$ ).

**Step 4:** Substitute back and solve for  $I$ :

$$2I = I - \frac{\pi}{2} \log 2 \implies I = -\frac{\pi}{2} \log 2$$

**Final Answer:**

$$I = -\frac{\pi}{2} \ln 2$$

**Quick Tip:** Use the King's rule (property P4) to relate  $I$  to  $\log \cos x$ , add, use  $\sin 2x$ , and self-reference the same integral.

34. Solve:  $\int \left[ \sqrt{\cot x} + \sqrt{\tan x} \right] dx.$

**Correct Answer:** —

**Solution:**

**Step 1:** Combine the two terms over a common form:

$$\sqrt{\cot x} + \sqrt{\tan x} = \frac{\cos x}{\sqrt{\sin x \cos x}} + \frac{\sin x}{\sqrt{\sin x \cos x}} = \frac{\sin x + \cos x}{\sqrt{\sin x \cos x}}$$

**Step 2:** Rewrite  $\sin x \cos x$  using the double angle identity:

$$\sin x \cos x = \frac{\sin 2x}{2}, \text{ so the integral becomes } \int \frac{\sqrt{2}(\sin x + \cos x)}{\sqrt{\sin 2x}} dx.$$

**Step 3:** Substitute  $t = \sin x - \cos x$ , so  $dt = (\cos x + \sin x) dx$ :

$$\text{Also } t^2 = (\sin x - \cos x)^2 = 1 - 2 \sin x \cos x = 1 - \sin 2x, \text{ so } \sin 2x = 1 - t^2.$$

The integral becomes  $\int \frac{\sqrt{2} dt}{\sqrt{1-t^2}}$ .

**Step 4:** Integrate using the standard arcsine form:

$$\sqrt{2} \int \frac{dt}{\sqrt{1-t^2}} = \sqrt{2} \sin^{-1}(t) + C$$

**Final Answer:**

$$\boxed{\sqrt{2} \sin^{-1}(\sin x - \cos x) + C}$$

**Quick Tip:** Combine both terms over  $\sqrt{\sin x \cos x}$ , substitute  $t = \sin x - \cos x$ , and integrate to an arcsine.