


SOLVED PAPER

VITEEE
2014

PART - I (PHYSICS)

- The amplification factor of a triode is 50. If the grid potential is decreased by 0.20 V. What increase, in plate potential will keep the plate current unchanged?
(a) 5 V (b) 10 V
(c) 0.2 V (d) 50 V
- If the nuclear fission, piece of uranium of mass 5.0 g is lost, the energy obtained in kWh is
(a) 1.25×10^7 (b) 2.25×10^7
(c) 3.25×10^7 (d) 0.25×10^7

- Current in the circuit will be



- $\frac{5}{40}$ A
 - $\frac{5}{50}$ A
 - $\frac{5}{10}$ A
 - $\frac{5}{20}$ A
- An installation consisting of an electric motor driving a water pump left 75 L of water per second to a height of 4.7 m. If the motor consumes a power of 5 kW, then the efficiency of the installation is
(a) 39% (b) 69%
(c) 93% (d) 96%
- A potential difference across the terminals of a battery is 50 V when 11 A current is drawn and 60 V, when 1 A current is drawn. The emf and the internal resistance of the battery are
(a) 62 V, 2 Ω (b) 63 V, 1 Ω
(c) 61 V, 1 Ω (d) 64 V, 2 Ω

- Beyond which frequency, the ionosphere bands any incident electromagnetic radiation but do not reflect it back towards the earth?
(a) 50 MHz (b) 40 MHz
(c) 30 MHz (d) 20 MHz
- A metallic surface ejects electrons. When exposed to green light of intensity I but no photoelectrons are emitted, when exposed to yellow light of intensity I. It is possible to eject electron from the same surface by
(a) yellow light of same intensity which is more than I
(b) green light of any intensity
(c) red light of any intensity
(d) None of the above
- An electron moves at right angle to a magnetic field of 5×10^{-2} T with a speed of 6×10^7 m/s. If the specific charge of the electron is 1.7×10^{11} C/kg. The radius of the circular path will be
(a) 2.9 cm (b) 3.9 cm
(c) 2.35 cm (d) 2 cm
- A solenoid 30 cm long is made by winding 2000 loops of wire on an iron rod whose cross-section is 1.5 cm^2 . If the relative permeability of the iron is 6000. What is the self-inductance of the solenoid?
(a) 1.5 H (b) 2.5 H
(c) 3.5 H (d) 0.5 H
- A coil of resistance 10 Ω and an inductance 5 H is connected to a 100 V battery. The energy stored in the coil is
(a) 325 erg (b) 125 J
(c) 250 erg (d) 250 J
- A galvanometer has current range of 15 mA and voltage range 750 mV. To convert this galvanometer into an ammeter of range 25 A, the required shunt is
(a) 0.8 Ω (b) 0.93 Ω
(c) 0.03 Ω (d) 2.0 Ω

12. The denial cell is balanced on 125 cm length of a potentiometer. Now, the cell is short circuited by a resistance of 2Ω and the balance is obtained at 100 cm. The internal resistance of the denial cell is

(a) $\frac{4}{3}\Omega$ (b) 1.5Ω
(c) 1.25Ω (d) 0.5Ω


13. Four resistance of 10Ω , 60Ω , 100Ω and 200Ω respectively taken in order are used to form a Wheatstone's bridge. A 15V battery is connected to the ends of a 200Ω resistance, the current through it will be

(a) $7.5 \times 10^{-5}\text{ A}$ (b) $7.5 \times 10^{-4}\text{ A}$
(c) $7.5 \times 10^{-3}\text{ A}$ (d) $7.5 \times 10^{-2}\text{ A}$

14. A circuit has a self-inductance of 1 H and carries a current of 2A. To prevent sparking, when the circuit is switched off, a capacitor which can withstand 400 V is used. The least capacitance of capacitor connected across the switch must be equal to

(a) $50\mu\text{F}$ (b) $25\mu\text{F}$
(c) $100\mu\text{F}$ (d) $12.5\mu\text{F}$

15. The output Y of the logic circuit shown in figure is best represented as



(a) $\overline{A} + \overline{B.C}$ (b) $A + \overline{B.C}$
(c) $\overline{A + B.C}$ (d) $\overline{A + \overline{B.C}}$

16. A resistor of $6\text{ k}\Omega$ with tolerance 10% and another resistance of $4\text{ k}\Omega$ with tolerance 10% are connected in series. The tolerance of the combination is about

(a) 5% (b) 10%
(c) 12% (d) 15%

17. If we add impurity to a metal those atoms also deflect electrons. Therefore,

(a) the electrical and thermal conductivities both increase
(b) the electrical and thermal conductivities both decrease
(c) the electrical conductivity increases but thermal conductivity decreases
(d) the electrical conductivity decrease but thermal conductivity increases

18. A proton and an α -particle, accelerated through the same potential difference, enter a region of uniform magnetic field normally. If the radius of the proton orbit is 10 cm, then radius of α -particle is

(a) 10 cm (b) $10\sqrt{2}\text{ cm}$
(c) 20 cm (d) $5\sqrt{2}\text{ cm}$

19. An ammeter and a voltmeter of resistance R are connected in series to an electric cell of negligible internal resistance. Their reading are A and V respectively. If another resistance R is connected in parallel with the voltmeter, then

(a) both A and V will increase
(b) both A and V will decrease
(c) A will decrease and V will increase
(d) A will increase and V will decrease

20. A neutron is moving with velocity u . It collides head on and elastically with an atom of mass number A . If the initial kinetic energy of the neutron is E , then how much kinetic energy will be retained by the neutron after reflection?

(a) $\left(\frac{A}{A+1}\right)^2 E$ (b) $\frac{A}{(A+1)^2} E$
(c) $\left(\frac{A-1}{A+1}\right)^2 E$ (d) $\frac{(A-1)}{(A+1)^2} E$


21. If a magnet is suspended at angle 30° to the magnet meridian, the dip of needle makes angle of 45° with the horizontal, the real dip is

(a) $\tan^{-1}\left(\frac{\sqrt{3}}{2}\right)$ (b) $\tan^{-1}(\sqrt{3})$
(c) $\tan^{-1}\left(\sqrt{\frac{3}{2}}\right)$ (d) $\tan^{-1}\left(\frac{2}{\sqrt{3}}\right)$

22. Which has more luminous efficiency?


(a) A 40 W bulb
(b) A 40W fluorescent tube
(c) Both have same
(d) Cannot say

23. The resistance of a germanium junction diode whose $V-I$ is shown in figure is ($V_k = 0.3\text{ V}$)



- (a) $5\text{ k}\Omega$ (b) $0.2\text{ k}\Omega$
 (c) $2.3\text{ k}\Omega$ (d) $\left(\frac{10}{2.3}\right)\text{ k}\Omega$


24. In hydrogen discharge tube, it is observed that through a given cross-section 3.31×10^{15} electrons are moving from right to left and 3.12×10^5 protons are moving from left to right. The current in the discharge tube and its direction will be
 (a) 2 mA towards left
 (b) 2 mA, towards right
 (c) 1 mA, towards right
 (d) 2 mA, towards left
25. In a semiconductor, separation between conduction and valence band is of the order of
 (a) 0 eV (b) 1 eV
 (c) 10 eV (d) 50 eV
26. If 1000 droplets each of potential 1 V and radius r are mixed to form a big drop. Then, the potential of the drop as compared to small droplets, will be
 (a) 1000 V (b) 800 V
 (c) 100 V (d) 20 V
27. A Zener diode, having breakdown voltage equal to 15 V is used in a voltage regulator circuit shown in figure. The current through the diode is



- (a) 10 mA (b) 15 mA
 (c) 20 mA (d) 5 mA
28. The activity of a radioactive sample is measured as N_0 counts per minute at $t = 0$ and N_0/C counts per minute at $t = 5$ min. The time, (in minute) at which the activity reduces to half its value, is
 (a) $\log_e \frac{2}{5}$ (b) $\frac{5}{\log_e 2}$
 (c) $5 \log_{10} 2$ (d) $5 \log_e 2$
29. If the electron in the hydrogen atom jumps from third orbit to second orbit, the wavelength of the emitted radiation in term of Rydberg constant is

- (a) $\frac{6}{5R}$ (b) $\frac{36}{5R}$
 (c) $\frac{64}{7R}$ (d) None of these

30. Silver has a work function of 4.7 eV. When ultraviolet light of wavelength 100 nm is incident on it a potential of 7.7 V is required to stop the photoelectrons from reaching the collector plate. How much potential will be required to stop photoelectrons, when light of wavelength 200 nm is incident on it?
 (a) 15.4 V (b) 2.35 V
 (c) 3.85 V (d) 1.5 V
31. If the distance of 100 W lamp is increased from a photocell, the saturation current i in the photocell varies with the distance d as
 (a) $i \propto d^2$ (b) $i \propto d$
 (c) $i \propto \frac{1}{d}$ (d) $i \propto \frac{1}{d^2}$
32. Following process is known as
 $h\nu \longleftrightarrow e^+ + e^-$
 (a) Pair production (b) photoelectric effect
 (c) Compton effect (d) Zeeman effect
33. During charging a capacitor, variations of potential V of the capacitor with time t is shown as



34. When a resistor of $11\ \Omega$ is connected in series with a electric cell. The current following in it is 0.5 A. Instead when a resistor of $5\ \Omega$ is connected to the same electric cell in series, the current increases by 0.4 A. The internal resistance of the cell is
 (a) $1.5\ \Omega$ (b) $2\ \Omega$
 (c) $2.5\ \Omega$ (d) $3.5\ \Omega$

35. A battery is charged at a potential of 15 V in 8 h when the current flowing is 10 A. The battery on discharge supplies a current of 5 A for 15 h. The mean terminal voltage during discharge is 14 V. The watt-hour efficiency of battery is
 (a) 80% (b) 90%
 (c) 87.5% (d) 82.5%
36. A circular current carrying coil has a radius R. The distance from the centre of the coil on the axis, where the magnetic induction will be $\frac{1}{8}$ th to its value at the centre of the coil is
 (a) $\frac{R}{\sqrt{3}}$ (b) $R\sqrt{3}$
 (c) $2\sqrt{3}R$ (d) $\frac{2}{\sqrt{3}}R$
37. The incorrect statement regarding the lines of force of the magnetic field B is
 (a) magnetic intensity is a measure of lines of force passing through unit area held normal to it
 (b) magnetic lines of force forms a close curve
 (c) inside a magnet, its magnetic lines of force move from north pole of a magnetic towards its south pole
 (d) due to a magnetic lines of force never cut each other
38. Two coils have a mutual inductance 0.55 H. The current changes in the first coil according to equation $I = I_0 \sin \omega t$. where, $I_0 = 10$ A and $\omega = 100\pi$ rad/s. The maximum value of emf in the second coil is
 (a) 2π (b) 5π
 (c) π (d) 4π
39. An L-C-R circuit contains $R = 50\ \Omega$, $L = 1$ mH and $C = 0.1\ \mu\text{F}$. The impedance of the circuit will be minimum for a frequency of
 (a) $\frac{10^5}{2\pi}$ Hz (b) $\frac{10^6}{2\pi}$ Hz
 (c) $2\pi \times 10^5$ Hz (d) $2\pi \times 10^6$ Hz
40. An eye can detect 5×10^4 photons per square meter per sec of green light ($\lambda = 5000\text{\AA}$) while the ear can detect 10^{-13} W/m^2 . The factor by which the eye is more sensitive as a power detector than ear is close to
 (a) 5 (b) 10
 (c) 10^6 (d) 15



PART - II(CHEMISTRY)

41. The sodium extract of an organic compound on acidification with acetic acid and addition of lead acetate solution gives a black precipitate. The organic compound contains
 (a) nitrogen (b) halogen
 (c) sulphur (d) phosphorus
42. The volume strength of 1.5 N H_2O_2 solution is
 (a) 16.8 L (b) 8.4 L
 (c) 4.2 L (d) 5.2 L
43. $\text{MnO}_4^- + 8\text{H}^+ + 5\text{e}^- \longrightarrow \text{Mn}^{2+} + 4\text{H}_2\text{O}$;
 $E^\circ = 1.51\text{ V}$
 $\text{MnO}_2 + 4\text{H}^+ + 2\text{e}^- \longrightarrow \text{Mn}^{2+} + 2\text{H}_2\text{O}$;
 $E^\circ = 1.23\text{ V}$ $E^\circ_{\text{MnO}_4^-/\text{MnO}_2}$ is
 (a) 1.70 V (b) 0.91 V
 (c) 1.37 V (d) 0.548 V
44. A metal has bcc structure and the edge length of its unit cell is 3.04\AA . The volume of the unit cell in cm^3 will be
 (a) $1.6 \times 10^{21}\text{ cm}^3$ (b) $2.81 \times 10^{-23}\text{ cm}^3$
 (c) $6.02 \times 10^{-23}\text{ cm}^3$ (d) $6.6 \times 10^{-24}\text{ cm}^3$
45. Among $[\text{Fe}(\text{H}_2\text{O})_6]^{3+}$, $[\text{Fe}(\text{CN})_6]^{3-}$, $[\text{Fe}(\text{Cl})_6]^{3-}$ species, the hybridisation state of the Fe atom are, respectively.
 (a) d^2sp^3 , d^2sp^3 , sp^3d^2 (b) sp^3d^2 , d^2sp^3 , d^2sp^3
 (c) sp^3d^2 , d^2sp^3 , sp^3d^2 (d) None of the above
46. Which of the following hydrogen bonds are strongest in vapour phase?
 (a) HF HF
 (b) HF HCl
 (c) HCl HCl
 (d) HF HI
47. The rate constant for forward reaction and backward reaction of hydrolysis of ester are 1.1×10^{-2} and 1.5×10^{-3} per minute respectively. Equilibrium constant for the reaction is
 $\text{CH}_3\text{COOC}_2\text{H}_5 + \text{H}_2\text{O} \rightleftharpoons \text{CH}_3\text{COOH} + \text{C}_2\text{H}_5\text{OH}$
 (a) 33.7 (b) 7.33
 (c) 5.33 (d) 33.3
48. 19.85 mL of 0.1 N NaOH reacts with 20 mL of HCl solution for complete neutralisation. The molarity of HCl solution is
 (a) 9.9 (b) 0.99
 (c) 0.099 (d) 0.0099

49. An *f*-shell containing 6 unpaired electrons can exchange
 (a) 6 electrons (b) 9 electrons
 (c) 12 electrons (d) 15 electrons
50. The standard molar heat of formation of ethane, CO₂ and water (l) are respectively -21.1, -94.1 and -68.3 kcal. The standard molar heat of combustion of ethane will be
 (a) -372 kcal (b) 162 kcal
 (c) -240 kcal (d) 183.5 kcal
51. The solubility product of Ag₂CrO₄ is 32×10^{-12} . What is the concentration of CrO₄⁻ ions in that solution?
 (a) 2×10^{-4} M (b) 16×10^{-4} M
 (c) 8×10^{-4} M (d) 8×10^{-8} M
52. The equivalent conductivity of a solution containing 2.54g of CuSO₄ per L is 91.0 W⁻¹ cm² eq⁻¹. Its conductivity would be
 (a) $2.9 \times 10^{-3} \Omega^{-1} \text{ cm}^{-1}$
 (b) $1.8 \times 10^{-2} \Omega^{-1} \text{ cm}^{-1}$
 (c) $2.4 \times 10^{-4} \Omega^{-1} \text{ cm}^{-1}$
 (d) $3.6 \times 10^{-3} \Omega^{-1} \text{ cm}^{-1}$
53. The half-life of two samples are 0.1 and 0.8 s. Their respective concentration are 400 and 50 respectively. The order of the reaction is
 (a) 0 (b) 2
 (c) 1 (d) 4
54. Which sequence of reactions shows correct chemical relation between sodium and its compounds?
 (a) $\text{Na} + \text{O}_2 \longrightarrow \text{Na}_2\text{O} \xrightarrow{\text{HCl(aq)}} \text{NaCl} \xrightarrow{\text{CO}_2} \text{Na}_2\text{CO}_3 \xrightarrow{\Delta} \text{Na}$
 (b) $\text{Na} \xrightarrow{\text{O}_2} \text{Na}_2\text{O} \xrightarrow{\text{H}_2\text{O}} \text{NaOH} \xrightarrow{\text{CO}_2} \text{Na}_2\text{CO}_3 \xrightarrow{\Delta} \text{Na}$
 (c) $\text{Na} + \text{H}_2\text{O} \longrightarrow \text{NaOH} \xrightarrow{\text{HCl}} \text{NaCl} \xrightarrow{\text{CO}_2} \text{Na}_2\text{CO}_3 \xrightarrow{\Delta} \text{Na}$
 (d) $\text{Na} + \text{H}_2\text{O} \longrightarrow \text{NaOH} \xrightarrow{\text{CO}_2} \text{Na}_2\text{CO}_3 \xrightarrow[\text{(molten)}]{\text{HCl}} \text{NaCl} \xrightarrow{\text{Electrolysis}} \text{Na}$
55. Purest form of iron is
 (a) pig iron (b) wrought iron
 (c) cast iron (d) steel
56. Which has the smallest size?
 (a) Na⁺ (b) Mg²⁺
 (c) Al³⁺ (d) P⁵⁺
57. In the reaction,
 $8\text{Al} + 3\text{Fe}_3\text{O}_4 \longrightarrow 4\text{Al}_2\text{O}_3 + 9\text{Fe}$
 the number of electrons transferred from the reductant to the oxidant is
 (a) 8 (b) 4
 (c) 16 (d) 24
58. The bond angles of NH₃, NH₄⁺ and NH₂⁻ are in the order
 (a) NH₂⁻ > NH₃ > NH₄⁺
 (b) NH₄⁺ > NH₃ > NH₂⁻
 (c) NH₃ > NH₂⁻ > NH₄⁺
 (d) NH > NH₄⁺ > NH₂⁻
59. A gaseous mixture containing He, CH₄ and SO₂ was allowed to effuse through a fine hole then find what molar ratio of gases coming out initially? (Given mixture contains He, CH₄ and SO₂ in 1 : 2 : 3 mole ratio).
 (a) $\sqrt{2} : \sqrt{2} : 3$ (b) 2 : 2 : 3
 (c) 4 : 4 : 3 (d) 1 : 1 : 3
60. According to Bohr's theory, the angular momentum for an electron of 3rd orbit is
 (a) $3\hbar$ (b) $1.5\hbar$
 (c) $9\hbar$ (d) $2\frac{\hbar}{\pi}$
61. 2.76 g of silver carbonate on being strongly heated yields a residue weighing
 (a) 3.54 g (b) 3.0 g
 (c) 1.36 g (d) 2.16 g
62. The final product (IV) in the sequence of reactions

$$\begin{array}{c} \text{CH}_3\text{CHOH} \xrightarrow{\text{PBr}_3} \text{I} \xrightarrow[\text{Ether}]{\text{Mg}} \\ | \\ \text{CH}_3 \end{array}$$

$$\begin{array}{c} \text{CH}_2\text{---CH}_2 \\ \diagup \quad \diagdown \\ \text{O} \\ \text{II} \end{array} \xrightarrow{\text{H}_2\text{O}} \text{III} \xrightarrow{\text{H}_2\text{O}} \text{IV is}$$
 (a) $\text{CH}_3\text{---CH}(\text{OCH}_2\text{CH}_2\text{OH})$
 (b) $\text{CH}_3\text{---CH}(\text{CH}_3)\text{CH}_2\text{CH}_2\text{Br}$
 (c) $\text{CH}_3\text{---CH}(\text{CH}_3)\text{---CH}_2\text{CH}_2\text{OH}$
 (d) $\text{CH}_3\text{---CH}(\text{CH}_3)\text{OCH}_2\text{CH}_3$



64. Which of the following has an ester linkage?

- (a) Nylon-66 (b) Dacron
(c) PVC (d) Bakelite

65. Which of the following pairs give positive Tollen's test?

- (a) Glucose, sucrose
(b) Glucose, fructose
(c) Hexanal, acetophenone
(d) Fructose, sucrose

66. Peptisation involves

- (a) precipitation of colloidal particles
(b) disintegration of colloidal aggregates
(c) evaporation of dispersion medium
(d) impact of molecules of the dispersion medium on the colloidal particles

67. Which of the following has the maximum number of unpaired d-electrons?

- (a) Fe^{2+} (b) Cu^+
(c) Zn (d) Ni^{3+}

68. Iodine is formed when potassium iodide reacts with a solution of

- (a) ZnSO_4 (b) CuSO_4
(c) $(\text{NH}_4)_2\text{SO}_4$ (d) Na_2SO_4

69. Which of the following does not represent the correct order of the property indicated?

- (a) $\text{Sc}^{3+} > \text{Cr}^{3+} > \text{Fe}^{3+} > \text{Mn}^{3+}$ — ionic radii
(b) $\text{Sc} < \text{Ti} < \text{Cr} < \text{Mn}$ — density
(c) $\text{Mn}^{2+} > \text{Ni}^{2+} > \text{Co}^{2+} < \text{Fe}^{2+}$ — ionic radii
(d) $\text{FeO} < \text{CaO} < \text{MnO} < \text{CuO}$ — basic nature

70. If the elevation in boiling point of a solution of 10 g of solute (mol. wt. = 100) in 100 g of water is ΔT_b , the ebullioscopic constant of water is

- (a) 10 (b) $100 T_b$
(c) $\frac{\Delta T_b}{10}$

71. Which of the following compounds cannot be prepared singly by the Wurtz reaction?

- (a) C_2H_6
(b) $(\text{CH}_3)_2\text{CHCH}_3$
(c) $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_3$
(d) All of the above can be prepared

72. Which of the following oxides is strongly basic?

- (a) Ti_2O (b) B_2O_3
(c) Al_2O_3 (d) Ga_2O_3





73. In Langmuir's model of adsorption of a gas on a solid surface,

- (a) the rate of dissociation of adsorbed molecules from the surface does not depend on the surface covered
(b) the adsorption at a single site on the surface may involve multiple molecules at the same time
(c) the mass of gas striking a given area of surface is proportional to the pressure of the gas
(d) the mass of gas striking a given area of surface is independent of the pressure of the gas


74. How many sigma and pi-bonds are there in the molecule of dicyanoethene ($\text{CN}-\text{CH}=\text{CH}-\text{CN}$)?

- (a) 3 sigma and 3 pi (b) 5 sigma and 2 pi
(c) 7 sigma and 5 pi (d) 2 sigma and 3 pi


75. What will be the order of reactivity of the following carbonyl compounds with Grignard's reagent?



- (a) $\text{I} > \text{II} > \text{III} > \text{IV}$ (b) $\text{IV} > \text{III} > \text{II} > \text{I}$
(c) $\text{II} > \text{I} > \text{IV} > \text{III}$ (d) $\text{III} > \text{II} > \text{I} > \text{IV}$





The final product 'C' in the above reaction is



77. Which of the following isomerism is shown by ethyl acetoacetate?

- (a) Geometrical isomerism
(b) Keto-enol tautomerism
(c) Enantiomerism
(d) Diastereoisomerism

78. The final product obtained in the reaction,



79. Among the following the strongest nucleophile is

- (a) C_2H_5SH (b) CH_3COO^-
(c) CH_3NH_2 (d) $NCCH_2$

80. Which set has different class of compounds?

- (a) Tranquillizers-Equanil, heroin, valium
(b) Antiseptics-Bithional, dettol, boric acid
(c) Analgesics-Naproxen, morphine, aspirin
(d) Bactericidal-penicillin, aminoglycosides, ofloxacin

PART-III(MATHEMATICS)

81. The solution of $\frac{dy}{dx} = \frac{x^2 + y^2 + 1}{2xy}$, satisfying

$y(1) = 0$ is given by

- (a) hyperbola (b) circle
(c) ellipse (d) parabola

82. If $x \cdot \frac{dy}{dx} + y = x \cdot \frac{f(xy)}{f'(xy)}$, then $f(xy)$ is equal to

- (a) $k \cdot e^{\frac{x^2}{2}}$ (b) $k \cdot e^{y^{2/2}}$
(c) $k \cdot e^{x^2}$ (d) $k \cdot e^{\frac{xy}{2}}$

83. The differential equation of the rectangular hyperbola hyperbola, where axes are the asymptotes of the hyperbola, is

- (a) $y \frac{dy}{dx} = x$ (b) $x \frac{dy}{dx} = -y$
(c) $x \frac{dy}{dx} = y$ (d) $xdy + ydx = c$

84. The length of longer diagonal of the parallelogram constructed on $5a + 2b$ and $a - 3b$, if it is given that $|a| = 2\sqrt{2}$, $|b| = 3$ and the angle

between a and b is $\frac{\pi}{4}$, is

- (a) 15 (b) $\sqrt{113}$
(c) $\sqrt{593}$ (d) $\sqrt{369}$

85. If $r = \alpha b \times c + \beta c \times a + \gamma a \times b$ and $[a \ b \ c] = 2$, then $\alpha + \beta + \gamma$ is equal to

- (a) $r \cdot [b \times c + c \times a + a \times b]$
(b) $\frac{1}{2} r \cdot (a + b + c)$
(c) $2r \cdot (a + b + c)$
(d) 4

86. If a, b, c are three non-coplanar vectors and p, q, r are reciprocal vectors, then $(la + mb + nc) \cdot (lp + mq + nr)$ is equal to

- (a) $l + m + n$ (b) $l^3 + m^3 + n^3$
(c) $l^2 + m^2 + n^2$ (d) None of these

87. If the integers m and n are chosen at random from 1 to 100, then the probability that a number of the form $7^n + 7^m$ is divisible by 5, equals to

- (a) $\frac{1}{4}$ (b) $\frac{1}{2}$ (c) $\frac{1}{8}$ (d) $\frac{1}{3}$

88. Let X denote the sum of the numbers obtained when two fair dice are rolled. The variance and standard deviation of X are

(a) $\frac{31}{6}$ and $\sqrt{\frac{31}{6}}$ (b) $\frac{35}{6}$ and $\sqrt{\frac{35}{6}}$

(c) $\frac{17}{6}$ and $\sqrt{\frac{17}{6}}$ (d) $\frac{31}{6}$ and $\sqrt{\frac{35}{6}}$

89. A four digit number is formed by the digits 1, 2, 3, 4 with no repetition. The probability that the number is odd, is

(a) zero (b) $\frac{1}{3}$

(c) $\frac{1}{4}$ (d) None of these

90. If the vertices of a triangle are $A(0, 4, 1)$, $B(2, 3, -1)$ and $C(4, 5, 0)$, then the orthocentre of $\triangle ABC$, is

(a) $(4, 5, 0)$ (b) $(2, 3, -1)$
(c) $(-2, 3, -1)$ (d) $(2, 0, 2)$

91. The equation of normal to the curve $y = (1+x)^y + \sin^{-1}(\sin^2 x)$ at $x = 0$ is

(a) $x + y = 1$ (b) $x - y = 1$
(c) $x + y = -1$ (d) $x - y = -1$

92. The value of c from the Lagrange's mean value theorem for which $f(x) = \sqrt{25 - x^2}$ in $[1, 5]$, is

(a) 5 (b) 1
(c) $\sqrt{15}$ (d) None of these

93. If $A = \begin{bmatrix} 3 & 4 \\ 5 & 7 \end{bmatrix}$, then $A \cdot (\text{adj } A)$ is equal to

(a) A (b) $|A|$
(c) $|A|$ (d) None of these

94. If there is an error of $k\%$ in measuring the edge of a cube, then the percent error in estimating its volume is

(a) k (b) $3k$
(c) $\frac{k}{3}$ (d) None of these

95. If the system of equations $x + ky - z = 0$, $3x - ky - z = 0$ and $x - 3y + z = 0$, has non-zero solution, then k is equal to

(a) -1 (b) 0
(c) 1 (d) 2

96. If the points $(1, 2, 3)$ and $(2, -1, 0)$ lie on the opposite sides of the plane $2x + 3y - 2z = k$, then

(a) $k < 1$ (b) $k > 2$
(c) $k < 1$ or $k > 2$ (d) $1 < k < 2$

97. If $\Delta(x) = \begin{vmatrix} 1 & \cos x & 1 - \cos x \\ 1 + \sin x & \cos x & 1 + \sin x - \cos x \\ \sin x & \sin x & 1 \end{vmatrix}$,

then $\int_0^{\pi/4} \Delta(x) dx$ is equal to

(a) $\frac{1}{4}$ (b) $\frac{1}{2}$ (c) 0 (d) $-\frac{1}{4}$

98. Let $f'(x)$, be differentiable $\forall x$. If $f(1) = -2$ and $f'(x) \geq 2 \forall x \in [1, 6]$, then

(a) $f(6) < 8$ (b) $f(6) \geq 8$
(c) $f(6) \geq 5$ (d) $f(6) \leq 5$

99. If $\Delta_r = \begin{vmatrix} 2r-1 & {}^m C_r & 1 \\ m^2-1 & 2^m & m+1 \\ \sin^2(m^2) & \sin^2(m) & \sin^2(m+1) \end{vmatrix}$, then

the value of $\sum_{r=0}^m \Delta_r$, is

(a) 1 (b) 0
(c) 2 (d) None of these

100. Two lines $\frac{x-1}{2} = \frac{y+1}{3} = \frac{z-1}{4}$ and

$\frac{x-3}{1} = \frac{y-k}{2} = z$ intersect at a point, if k is equal to

(a) $\frac{2}{9}$ (b) $\frac{1}{2}$ (c) $\frac{9}{2}$ (d) $\frac{1}{6}$

101. The minimum value of $\frac{x}{\log x}$ is

(a) e (b) $\frac{1}{e}$
(c) e^2 (d) e^3

102. The triangle formed by the tangent to the curve $f(x) = x^2 + bx - b$ at the point $(1, 1)$ and the coordinate axes lies in the first quadrant. If its area is 2, then the value of b is

(a) -1 (b) 3
(c) -3 (d) 1

103. The statement $(p \Rightarrow q) \Leftrightarrow (\sim p \wedge q)$ is a

(a) tautology
(b) contradiction
(c) Neither (a) nor (b)
(d) None of these

104. If $x + iy = \frac{3}{2 + \cos \theta + i \sin \theta}$, then $x^2 + y^2$ is equal to

(a) $3x - 4$ (b) $4x - 3$
(c) $4x + 3$ (d) None of these

105. The negation of $(\sim p \wedge q) \vee (p \wedge \sim q)$ is
 (a) $(p \vee \sim q) \vee (\sim p \vee q)$
 (b) $(p \vee \sim q) \wedge (\sim p \vee q)$
 (c) $(p \wedge \sim q) \wedge (\sim p \vee q)$
 (d) $(p \wedge \sim q) \wedge (p \vee \sim q)$
106. The normals at three points P, Q and R of the parabola $y^2 = 4ax$ meet at (h, k) . The centroid of the $\triangle PQR$ lies on
 (a) $x=0$ (b) $y=0$
 (c) $x=-a$ (d) $y=a$
107. The minimum area of the triangle formed by any tangent to the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ with the coordinate axes is
 (a) $a^2 + b^2$ (b) $\frac{(a+b)^2}{2}$
 (c) ab (d) $\frac{(a-b)^2}{2}$
108. If the line $lx + my - n = 0$ will be a normal to the hyperbola, then $\frac{a^2}{l^2} - \frac{b^2}{m^2} = \frac{(a^2 + b^2)^2}{k}$, where k is equal to
 (a) n (b) n^2
 (c) n^3 (d) None of these
109. If $\cos \alpha + i \sin \alpha, b = \cos \beta + i \sin \beta$,
 $c = \cos \gamma + i \sin \gamma$ and $\frac{b}{c} + \frac{c}{a} + \frac{a}{b} = 1$, then $\cos(\beta - \gamma) + \cos(\gamma - \alpha) + \cos(\alpha - \beta)$ is equal to
 (a) $\frac{3}{2}$ (b) $-\frac{3}{2}$
 (c) 0 (d) 1
110. If $|z + 4| \leq 3$, then the greatest and the least value of $|z + 1|$ are
 (a) $-1, 6$ (b) $6, 0$
 (c) $6, 3$ (d) None of these
111. The angle between lines joining the origin to the point of intersection of the line $\sqrt{3}x + y = 2$ and the curve $y^2 - x^2 = 4$ is
 (a) $\tan^{-1} \frac{2}{\sqrt{3}}$ (b) $\frac{\pi}{6}$
 (c) $\tan^{-1} \left(\frac{\sqrt{3}}{2} \right)$ (d) $\frac{\pi}{2}$
112. If the area of the triangle on the complex plane formed by the points $z, z + iz$ and iz is 200, then the value of $3|z|$ must be equal to
 (a) 20 (b) 40
 (c) 60 (d) 80
113. Equation of the chord of the hyperbola $25x^2 - 16y^2 = 400$ which is bisected at the point $(6, 2)$ is
 (a) $6x - 7y = 418$ (b) $75x - 16y = 418$
 (c) $25x - 4y = 400$ (d) None of these
114. If a plane meets the coordinate axes at A, B and C such that the centroid of the triangle is $(1, 2, 4)$, then the equation of the plane is
 (a) $x + 2y + 4z = 12$ (b) $4x + 2y + z = 12$
 (c) $x + 2y + 4z = 3$ (d) $4x + 2y + z = 3$
115. The volume of the tetrahedron included between the plane $3x + 4y - 5z - 60 = 0$ and the coordinate planes is
 (a) 60 (b) 600
 (c) 720 (d) 400
116. $\int_0^{2\pi} (\sin x + |\sin x|) dx$ is equal to
 (a) 0 (b) 4
 (c) 8 (d) 1
117. The value of $\int_0^{\sqrt{2}} [x^2] dx$, where $[\cdot]$ is the greatest integer function, is
 (a) $2 - \sqrt{2}$ (b) $2 + \sqrt{2}$
 (c) $\sqrt{2} - 1$ (d) $\sqrt{2} - 2$
118. If $l(m, n) = \int_0^1 t^m (1+t)^n dt$, then the expression for $l(m, n)$ in terms of $l(m+1, n+1)$ is
 (a) $\frac{2^n}{m+1} - \frac{n}{m+1} \cdot l(m+1, n-1)$
 (b) $\frac{n}{m+1} \cdot l(m+1, n-1)$
 (c) $\frac{2n}{m+1} + \frac{n}{m+1} \cdot l(m+1, n-1)$
 (d) $\frac{m}{n+1} \cdot l(m+1, n-1)$
119. The area in the first quadrant between $x^2 + y^2 = \pi^2$ and $y = \sin x$ is
 (a) $\frac{\pi^3 - 8}{4}$ (b) $\frac{\pi^3}{4}$
 (c) $\frac{\pi^3 - 16}{4}$ (d) $\frac{\pi^3 - 8}{2}$
120. The area bounded by $y = xe^{|x|}$ and lines $|x| = 1, y = 0$ is
 (a) 4 sq units (b) 6 sq units
 (c) 1 sq unit (d) 2 sq units