CBSE Class 12 Mathematics 2014 - Set 65/3

Question Paper

Conducted by the Central Board of Secondary Education (CBSE)

Time Allowed: 3 Hours **Maximum Marks: 100**

General Instructions

(i) All questions are compulsory.

(ii) The question paper consists of 29 questions divided into three sections A, B and C. Section A comprises of 10 questions of one mark each, Section B comprises of 12 questions of four marks each and Section C comprises of 7 questions of six marks each.

(iii) All questions in Section A are to be answered in one word, one sentence or as per the exact requirement of the question.

(iv) There is no overall choice. However, internal choice has been provided in 4 questions of four marks each and 2 questions of six marks each. You have to attempt only one of the alternatives in all such questions.

(v) Use of calculators is not permitted. You may ask for logarithmic tables, if required.

Section A

1. If A is a square matrix such that $A^2 = A$, then write the value of $7A - (I + A)^3$, where I is an identity matrix.

2. If $\begin{bmatrix} x-y & z \\ 2x-y & w \end{bmatrix} = \begin{bmatrix} -1 & 4 \\ 0 & 5 \end{bmatrix}$, find the value of $x + y$.

3. If $\tan^{-1} x + \tan^{-1} y = \frac{\pi}{4}$; $xy < 1$, then write the value of $x + y + xy$.

4. If $\begin{vmatrix} 3x & 7 \\ -2 & 4 \end{vmatrix} = \begin{vmatrix} 8 & 7 \\ 6 & 4 \end{vmatrix}$, find the value of x.

5. If $f(x) = \int_0^x t \sin t \, dt$, then write the value of $f'(x)$.

6. Find the value of 'p' for which the vectors $3\hat{i} + 2\hat{j} + 9\hat{k}$ and $\hat{i} - 2p\hat{j} + 3\hat{k}$ are parallel.

7. If $R = \{(x, y) : x + 2y = 8\}$ is a relation on $\mathbb{N}$, write the range of R.

8. If the cartesian equations of a line are $\frac{3-x}{5} = \frac{y+4}{7} = \frac{2z-6}{4}$, write the vector equation for the line.

9. If $\int_0^a \frac{1}{4 + x^2} \, dx = \frac{\pi}{8}$, find the value of a.

10. If $\vec{a}$ and $\vec{b}$ are perpendicular vectors, $|\vec{a} + \vec{b}| = 13$ and $|\vec{a}| = 5$, find the value of $|\vec{b}|$.

Section B

11. Solve the differential equation $(1+x^2)\frac{dy}{dx} + y = e^{\tan^{-1}x}$.

12(a). Show that the four points A, B, C and D with position vectors $4\hat{i}+5\hat{j}+\hat{k}$, $-\hat{j}-\hat{k}$, $3\hat{i}+9\hat{j}+4\hat{k}$ and $4(-\hat{i}+\hat{j}+\hat{k})$ respectively are coplanar.

OR

12(b). The scalar product of the vector $\vec{a} = \hat{i}+\hat{j}+\hat{k}$ with a unit vector along the sum of vectors $\vec{b} = 2\hat{i}+4\hat{j}-5\hat{k}$ and $\vec{c} = \lambda\hat{i}+2\hat{j}+3\hat{k}$ is equal to one. Find the value of λ and hence find the unit vector along $\vec{b}+\vec{c}$.

13(a). Evaluate: $\int_0^{\pi} \frac{4x\sin x}{1+\cos^2 x}\,dx$.

OR

13(b). Evaluate: $\int \frac{x+2}{\sqrt{x^2+5x+6}}\,dx$.

14(a). Find the value(s) of x for which $y = [x(x-2)]^2$ is an increasing function.

OR

14(b). Find the equations of the tangent and normal to the curve $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ at the point $(\sqrt{2}a, b)$.

15. If the function $f : \mathbb{R} \to \mathbb{R}$ be given by $f(x) = x^2+2$ and $g : \mathbb{R} \to \mathbb{R}$ be given by $g(x) = \frac{x}{x-1}$, $x \neq 1$, find $f \circ g$ and $g \circ f$ and hence find $f \circ g(2)$ and $g \circ f(-3)$.

16(a). Prove that: $\tan^{-1}\left[\frac{\sqrt{1+x}-\sqrt{1-x}}{\sqrt{1+x}+\sqrt{1-x}}\right] = \frac{\pi}{4} - \frac{1}{2}\cos^{-1}x$, $-\frac{1}{\sqrt{2}} \leq x \leq 1$.

OR

16(b). If $\tan^{-1}\left(\frac{x-2}{x-4}\right) + \tan^{-1}\left(\frac{x+2}{x+4}\right) = \frac{\pi}{4}$, find the value of x.

17. An experiment succeeds thrice as often as it fails. Find the probability that in the next five trials, there will be at least 3 successes.

18. If $y = Pe^{ax} + Qe^{bx}$, show that $\frac{d^2y}{dx^2} - (a+b)\frac{dy}{dx} + aby = 0$.

19. Using properties of determinants, prove that: $\begin{vmatrix} 1+a & 1 & 1 \\ 1 & 1+b & 1 \\ 1 & 1 & 1+c \end{vmatrix} = abc+bc+ca+ab$.

20. If $x = \cos t\,(3-2\cos^2 t)$ and $y = \sin t\,(3-2\sin^2 t)$, find the value of $\frac{dy}{dx}$ at $t = \frac{\pi}{4}$.

21. Find the particular solution of the differential equation $\log\left(\frac{dy}{dx}\right) = 3x+4y$, given that $y = 0$ when $x = 0$.

22. Find the value of p, so that the lines l_1 : $\frac{1-x}{3} = \frac{7y-14}{p} = \frac{z-3}{2}$ and l_2 : $\frac{7-7x}{3p} = \frac{y-5}{1} = \frac{6-z}{5}$ are perpendicular to each other. Also find the equations of a line passing through the point $(3, 2, -4)$ and parallel to line l_1.

Section C

23(a). Find the equation of the plane through the line of intersection of the planes $x + y + z = 1$ and $2x + 3y + 4z = 5$ which is perpendicular to the plane $x - y + z = 0$. Also find the distance of the plane obtained above, from the origin.

OR

23(b). Find the distance of the point $(2, 12, 5)$ from the point of intersection of the line $\vec{r} = 2\hat{i} - 4\hat{j} + 2\hat{k} + \lambda(3\hat{i} + 4\hat{j} + 2\hat{k})$ and the plane $\vec{r} \cdot (\hat{i} - 2\hat{j} + \hat{k}) = 0$.

24. Using integration, find the area of the region bounded by the triangle whose vertices are $(-1, 2)$, $(1, 5)$ and $(3, 4)$.

25. A manufacturing company makes two types of teaching aids A and B of Mathematics for class XII. Each type of A requires 9 labour hours of fabricating and 1 labour hour for finishing. Each type of B requires 12 labour hours for fabricating and 3 labour hours for finishing. For fabricating and finishing, the maximum labour hours available per week are 180 and 30 respectively. The company makes a profit of ₹ 80 on each piece of type A and ₹ 120 on each piece of type B. How many pieces of type A and type B should be manufactured per week to get a maximum profit? Make it as an LPP and solve it graphically. What is the maximum profit per week?

26(a). There are three coins. One is a two-headed coin (having heads on both faces), another is a biased coin that comes up heads 75% of the times and third is a biased coin that comes up tails 40% of the times. One of the three coins is chosen at random and tossed, and it shows heads. What is the probability that it was the two-headed coin?

OR

26(b). Two numbers are selected at random (without replacement) from the first six positive integers. Let X denote the larger of the two numbers obtained. Find the probability distribution of the random variable X, and hence find the mean of the distribution.

27. Two schools A and B want to award their selected students on the values of sincerity, truthfulness and helpfulness. The school A wants to award ₹ x each, ₹ y each and ₹ z each for the three respective values to 3, 2 and 1 students respectively with a total award money of ₹ $1,600$. School B wants to spend ₹ $2,300$ to award its 4, 1 and 3 students on the respective values (by giving the same award money to the three values as before). If the total amount of award for one prize on each value is ₹ 900, using matrices, find the award money for each value. Apart from these three values, suggest one more value which should be considered for award.

28. If the sum of the lengths of the hypotenuse and a side of a right triangle is given, show that the area of the triangle is maximum when the angle between them is 60°.

29. Evaluate: $\int \frac{1}{\sin^4 x + \sin^2 x \cos^2 x + \cos^4 x}\, dx.$

- End of Question Paper -