

Question Paper

Conducted by the Central Board of Secondary Education (CBSE)



Time Allowed: 2 Hours

Maximum Marks: 35

General Instructions

- (i) This question paper contains **12 questions**. All questions are compulsory.
- (ii) Section A: Questions 1 to 3 are very short answer questions, carrying 2 marks each.
- (iii) Section B: Questions 4 to 11 are short answer questions, carrying 3 marks each.
- (iv) Section C: Questions 12 are case-based questions, carrying 5 marks each.
- (v) Attempt one alternative where an internal choice is provided. Follow the instructions within each question.
- (vi) Use of calculators is not allowed.

Section A


1. How will you carry out the following conversions : (Any two)

(i) Propanal to Propane

(ii) Ethanal to But-2-enal

(iii) Ethanoic acid to ethanamide

2. In the given reaction



the rate of formation of NH_3 is $3.6 \times 10^{-4} \text{ mol L}^{-1} \text{ s}^{-1}$.

Calculate the

(i) rate of reaction, and


(ii) rate of disappearance of $\text{H}_2(\text{g})$.

3. (i) State Kohlrausch's law of independent migration of ions.

(ii) Calculate the degree of dissociation (α) of CH_3COOH if Λ_m and Λ_m^0 of CH_3COOH are $48 \text{ S cm}^2 \text{ mol}^{-1}$ and $400 \text{ S cm}^2 \text{ mol}^{-1}$ respectively.

Section B


4. (a) Write the hybridisation and magnetic character of the following complexes :



[Atomic number : Ni = 28, Co = 27, Fe = 26]

OR

(b) (i) Write the IUPAC name of the following complex :



(ii) What is the difference between an Ambidentate ligand and a Bidentate ligand?

(iii) Out of $[\text{Fe}(\text{NH}_3)_6]^{3+}$ and $[\text{Fe}(\text{C}_2\text{O}_4)_3]^{3-}$, which complex is more stable and why?



5. Calculate the emf of the following cell :



Given : $E^\circ_{\text{Zn}^{2+}/\text{Zn}} = -0.76 \text{ V}$ and $E^\circ_{\text{Ag}^+/\text{Ag}} = +0.80 \text{ V}$

$[\log 2 = 0.3010, \log 3 = 0.4771, \log 10 = 1]$

6. (a) Write the major products in the following :



OR

(b) (i) Oxidation of propanal is easier than propanone. Why?

(ii) How can you distinguish between Acetophenone and Benzophenone?

(iii) Draw the structure of the following derivative : 2,4-Dinitrophenylhydrazone of Propanone.

7. A first order reaction takes 30 minutes for 75% decomposition. Calculate $t_{1/2}$.

Given : $[\log 2 = 0.3, \log 3 = 0.48, \log 4 = 0.6, \log 5 = 0.7]$

8. (a) Write three differences between Lyophobic sol and Lyophilic sol.

OR

(b) (i) Write the dispersed phase and dispersion medium of 'milk'.

(ii) What is the cause of Brownian movement in colloidal particles?

(iii) Why does physisorption decrease with increase in temperature?

9.

$E^\ominus_{M^{2+}/M}$	Cr	Mn	Fe	Co	Ni	Cu	Zn
	-0.91	-1.18	-0.44	-0.28	-0.25	+0.34	-0.76

From the given $E^\ominus$ values of the first row transition elements, answer the following questions :

(i) Why is $E^\ominus_{Mn^{2+}/Mn}$ value highly negative as compared to other elements?

(ii) What is the reason for the irregularity in the above $E^\ominus$ values?

(iii) Why is $E^\ominus_{Cu^{2+}/Cu}$ value exceptionally positive?

10. (a) Write the equation involved in the following reactions :

(i) Rosenmund reduction

(ii) Etard reaction


(iii) Stephen reaction

OR

(b) (i) Which acid of each pair would you expect to be stronger? Give reason.

(I) CH_3COOH or $F-CH_2-COOH$

(II)



(ii) Distinguish between Pentan-2-one and Pentan-3-one.

11. Account for the following :

(i) Transition metals and their compounds show catalytic activities.

(ii) Zn, Cd and Hg are non-transition elements.


(iii) Zr and Hf are of almost identical atomic radii.

12. Read the following passage and answer the questions that follow :

Amines constitute an important class of organic compounds derived by replacing one or more hydrogen atoms of ammonia molecule by alkyl/aryl groups. Amines are usually formed from nitro compounds, halides, amides, etc. They exhibit hydrogen bonding which influences their physical properties. Alkyl amines are found to be stronger bases than ammonia. In aromatic amines, electron releasing and withdrawing groups, respectively increase and decrease their basic character. Reactions of amines are governed by availability of the unshared pair of electrons on nitrogen. Influence of the number of hydrogen atoms at nitrogen atom on the type of reactions and nature of products is responsible for identification and distinction between primary, secondary and tertiary amines. Reactivity of aromatic amines can be controlled by acylation process.



(i) Why does aniline not give Friedel-Crafts reaction?

(ii) Arrange the following in the increasing order of their pK_b values :



(iii) How can you distinguish between $CH_3CH_2NH_2$ and $(CH_3CH_2)_2NH$ by Hinsberg test?

(iv) (a) Write the structures of A and B in the following reactions :



OR

(b) How will you convert the following :

(I) Benzoic acid to aniline

(II) Aniline to p-bromoaniline

- End of Question Paper -